

## Catalogue of American Amphibians and Reptiles.

McCoy, C.J., and R.C. Vogt. 1994. *Graptemys*.

***Graptemys* Agassiz**  
**Map Turtles**

*Graptemys* Agassiz, 1857:436. Type-species, *Graptemys geographica* (Le Sueur, 1817) [= *Testudo geographica* Le Sueur, 1817:86], subsequent designation by Brown, 1908:114 (Bour and Dubois, 1983:42).

*Malacoclemmys*: Cope, 1875:53 (part).

*Malaclemys*: McDowell, 1964:274.

• **Content.** Twelve species are recognized: *Graptemys barbouri*, *G. ernsti*, *G. gibbonsi*, *G. pulchra* (*pulchra* group); *G. geographica* (*geographica* group); *G. caglei*, *G. ouachitensis*, *G. pseudogeographica*, *G. versa* (*pseudogeographica* group); *G. flavimaculata*, *G. nigrinoda*, *G. oculifera* (*oculifera* group).

• **Definition.** Turtles in the genus *Graptemys* are small to large aquatic emydine turtles, maximum adult size ranging from 15 cm carapace length (male *G. nigrinoda*) to 32 cm (female *G. barbouri*). Adult females are larger than males, and sexual dimorphism in size is extreme (maximum SDI 2.58, in *G. pulchra*; Gibbons and Lovich, 1990). Adult females of some species have grotesquely enlarged heads and massive jaws. The carapace outline is oval to "teardrop shaped" and usually elevated in profile. Vertebral spines, present in most species, vary from extreme knob-like or blade-shaped projections to a slightly serrate, elevated vertebral profile. Vertebral spines


are more prominent in juveniles and males. The posterior edge of the carapace is strongly serrate, and the carapacial scutes are smooth. The neural bones are hexagonal and short-sided anteriorly. The plastron is large, solid (not hinged), and has large bridges with weak buttresses. The entoplastron lies anterior to the humero-pectoral seam.

The ground color of the carapace is a shade of green or greenish brown with a pattern of lighter (light green, yellow, or orange) and sometimes darker (brown or black) circles, spots, or lines. The pattern frequently consists of a light reticulum of interconnecting lines on a darker background, hence the common name "map turtles," in allusion to the map-like pattern. The plastron is light yellow or tan with a dark pattern of either concentric rings or lines along the scute seams. The ground color of head, legs, and tail is dark green to black with a multilineate pattern of lighter color (green or yellow). The head pattern consists of a combination of spots or blotches above, behind, and below the orbits, with a series of temporal lines extending onto the neck. Each species, although variable, has a modal head pattern that is species-specific and which may function in species recognition during courtship (Vogt, 1993).

The toes are fully webbed, and the hind feet are much enlarged for swimming. Each foot has five claws, and the foreclaws in adult males of some species are elongate.

The skull usually is narrow to moderately broad in males and broader in females. The temporal arch is complete and the orbitonasal foramen is enlarged. The crushing surfaces of the jaws are not ridged, and are greatly broadened in adult females of some species. The anterior border of the inferior process of the parietal is thickened, and the pterygoid touches the exoccipital.

The karyotype is either  $2n = 50$  or  $2n = 52$  (Bickham and Carr, 1983).



**Map.** Present range of the genus *Graptemys*. The stars mark extralimital fossil localities; see species accounts for other fossil records.

• **Descriptions and Illustrations.** References to external features are listed in the species accounts, and only selected references to descriptions or illustrations follow: general descriptions with illustrations (Carr, 1952; Ernst and Barbour, 1972, 1989; Vogt, 1981; Conant and Collins, 1991); illustrated descriptions of species, *G. barbouri* (Carr and Marchand, 1942; Cagle, 1952); *G. caglei* (Haynes and McKown, 1974); *G. ernsti* (Mount, 1975; Shealey, 1976; Lovich and McCoy, 1992); *G. flavimaculata* (Cagle, 1954); *G. geographica* (Cahn, 1937; Carr, 1952; Vogt, 1981); *G. gibbonsi* (Dundee and Rossman, 1989; Lovich and McCoy, 1992); *G. nigrinoda* (Cagle, 1954; Folkerts and Mount, 1969; Mount, 1975); *G. oculifera* (Cagle, 1953a; Dundee and Rossman, 1989); *G. ouachitensis* (Cagle, 1953b; Vogt, 1981, 1993); *G. pseudogeographica* (Vogt, 1981, 1993); *G. pulchra* (Cagle, 1952; Mount, 1975; Shealey, 1976; Lovich and McCoy, 1992); *G. versa* (Smith and Sanders, 1952; Haynes and McKown, 1974; Vogt, 1993). Skull and skeletal characters are described by McDowell (1964), Ernst and Barbour (1972), Gaffney (1979), and Dobie (1981); and karyotypes by McKown (1972), Stock (1972), and Killebrew (1977).

• **Distribution.** The genus *Graptemys* is found in temperate North America from southwestern Quebec and southern Ontario, including Lake George and Lake Champlain, westward through the Great Lakes and Mississippi drainages to the upper Missouri River in North Dakota. Populations occur throughout the Mississippi Basin, westward to west-central Texas (Brazos River), and eastward to extreme southwestern Virginia (Tennessee River drainage). Species of *Graptemys* also occur in Gulf Coast drainages from the Florida Panhandle to Louisiana, including the Apalachicola Bay drainage, Escambia Bay drainage, Mobile Bay drainage, Pascagoula River system, and the Pearl River. West of the Mississippi Embayment along the Gulf Coast, *Graptemys* species occur in the Colorado River (Texas) and the Guadalupe-San Antonio River system. Isolated populations occur in the Susquehanna River drainage in Pennsylvania and Maryland, in the Delaware River from its mouth to Sussex County, New Jersey, and in the lower Hudson River, New York. Species distributions are mapped in Iverson (1992).

• **Fossil Record.** All known fossil *Graptemys* have been referred to Recent species, and most are from sites within the modern range of the genus. Jackson (1975), however, described fossil *Graptemys barbouri* from a site (Pleistocene, Rancholabrean) on the Santa Fe River, Gilchrist County, Florida, 200 km east of the present range of the species. Fossil *Graptemys geographica* have been found in Pleistocene deposits in Texas, Kansas, Tennessee, and Virginia (Preston, 1979; McCoy and Vogt, 1990). Fossil *G. pseudogeographica* have been reported from Pleistocene sites in Michigan (Wilson and Zug, 1966), Kansas (Preston, 1979), and Texas (Preston, 1979; Slaughter et al., 1962; Stovall and McAnulty, 1950). The Michigan material may actually be *G. geographica* rather than *G. pseudogeographica* (Holman, 1988).

• **Pertinent Literature.** Most of the pertinent literature on the genus was reviewed in the species accounts. A recent general review, with species descriptions, key, and illustrations was that of Ernst and Barbour (1989). Recent taxonomic reviews are Lovich and McCoy (1992, *G. pulchra* group) and Vogt (1993, *G. pseudogeographica* group). Avise et al. (1992) and Lamb et al. (1994) reported on mitochondrial DNA in *Graptemys*. Iverson (1992) provided a distributional review with range maps for all species. A hypothesis for phylogenetic position of the genus was provided by Seidel and Jackson (1990) and a preliminary phylogeny for the species was presented by Iverson (1992).

• **Key to Species.** The Catalogue account numbers (when available) are given in parentheses after the species name.

1. a. Vertebral keel low, without prominent spines or knobs ..... 2
- b. Vertebral keel with prominent spines or knobs ..... 3
2. a. Horizontal J-shaped postorbital spot; carapace scutes convex; maximum CL 183 mm ..... *G. versa* (280)
- b. Postorbital spot oval or triangular; carapace scutes smooth; maximum CL 273 mm ..... *G. geographica* (484)
3. a. Vertebral keel with blunt, rounded black knobs .....

- ..... *G. nigrinoda* (396)
- b. Vertebral keel without blunt, rounded black knobs ..... 4
4. a. Circular yellow or orange blotch on each pleural scute ... .. *G. flavimaculata* (403)
- b. Pleural scute marks not solid circular blotches ..... 5
5. a. Pleural scutes with yellow-orange rings ..... .. *G. oculifera* (422)
- b. Pleural scutes without yellow-orange ring markings .... 6
6. a. Interorbital blotch present ..... 7
- b. Interorbital blotch absent ..... 10
7. a. Interorbital and postorbital blotches fused, with a central dark heart-shaped figure; chin with transverse light bar, rostrum blunt ..... *G. barbouri* (421)
- b. Interorbital and postorbital blotches fused or separate; chin with longitudinal light stripe; rostrum pointed ..... 8
8. a. Interorbital blotch not connected to postorbital blotches; nasal trident well-developed ..... *G. ernsti* (585)
- b. Interorbital and postorbital blotches connected; nasal trident present or absent ..... 9
9. a. Nasal trident usually present; single wide yellow bar (16-20% scute width) on upper side of each marginal scute .. .. *G. gibbonsi* (586)
- b. Nasal trident usually absent; narrow concentric yellow ocelli on marginal scutes ..... *G. pulchra* (360, part)
10. a. Chin with alternating transverse cream and dark green bars ..... 11
- b. Chin without transverse cream and dark green bars ... 12
11. a. Postorbital blotch chevron-shaped, usually connected to suborbital spot to form a crescent ..... *G. caglei* (184)
- b. Postorbital blotch reduced to a spot that is encircled with one or more narrow rings and never connected to suborbital spot to form a crescent ..... *G. ouachitensis sabinensis*
12. a. Postorbital blotch narrow or a narrow crescent; up to 6 lines entering orbit; no large spots on lower jaw ..... 13
- b. Postorbital blotch wide, or connected to subocular spot to form wide crescent; four large spots on lower jaw (mandibles, symphysis, throat) . *G. ouachitensis ouachitensis*
13. a. Postorbital blotch usually connected to subocular spot to form crescent, or no more than 3 lines entering orbit ..... .. *G. pseudogeographica kobnii*
- b. Postorbital blotch never connected to subocular spot; usually 3 or more (1-6) lines entering orbit ..... .. *G. pseudogeographica pseudogeographica*

• **Etymology.** The generic name *Graptemys* is derived from the Greek *graptos*, meaning inscribed or painted, and the Greek *emydos*, a freshwater turtle, in allusion to the often striking and complex color pattern of the carapace found in many species. The gender is feminine.

• **Comment.** The original concept of the genus *Graptemys* (Agassiz, 1857), as distinct from *Malaclemys*, was not immediately accepted. Neither Cope (1875) nor Boulenger (1889), in their influential catalogues, recognized *Graptemys* and the name *Malaclemys* (or *Malacoclemmys*, an invalid emendation) was generally used for species of *Graptemys* through the end of the 19th century. The type-species of *Graptemys* (*Testudo geographica* Le Sueur, 1817) was designated by Brown (1908), and the current concept of the genus was reestablished by Stejneger and Barbour (1917). That checklist set the pattern for the subsequent 47 years, until McDowell (1964) again proposed uniting *Graptemys* and *Malaclemys*. McDowell's proposal has not been widely accepted, notwithstanding that his arrangement has yet to be subjected to rigorous analysis. However consensus, supported by fragmentary evidence such as that supplied by McKown (1972), Wood (1977), and Dobie (1981), maintains that the two genera, however closely related, are on separate evolutionary trajectories.

## Literature Cited

- Agassiz, L. 1857. Contributions to the natural history of the United States. Vol. 1. Little, Brown and Co., Boston.
- Avise, J.C., B.W. Bowen, T. Lamb, A.B. Meylan, and E. Bermingham. 1992. Mitochondrial DNA evolution at a turtle's pace: evidence for low genetic variability and reduced microevolutionary rate in the Testudines. *Mol. Biol. Evol.* 9:457-473.
- Bickham, J.W. and J.L. Carr. 1983. Taxonomy and phylogeny of the higher categories of cryptodiran turtles based on a cladistic analysis of chromosomal data. *Copeia* 1983:918-932.
- Boulenger, G.A. 1889. Catalogue of the Chelonians, Rhynchocephalians, and Crocodiles in the British Museum (Natural History). The Trustees (Brit. Mus. Nat. Hist.), London.
- Bour, R. and A. Dubois. 1983. Statut nomenclatural et specimens-types d'*Emys pseudogeographica* Gray, 1831 et d'*Emys lesueuri* Gray, 1831 (Reptilia, Chelonii, Emydidae). *Bull. Mens. Soc. Linnéenne Lyon* 52:42-46.
- Brown, A.E. 1908. Generic types of Nearctic Reptilia and Amphibia. *Proc. Acad. Nat. Sci. Philadelphia* 60:112-127.
- Cagle, F.R. 1952. The status of the turtles *Graptemys pulchra* Baur and *Graptemys barbouri* Carr and Marchand, with notes on their natural history. *Copeia* 1952:223-234.
- . 1953a. The status of the turtle *Graptemys oculifera* (Baur). *Zoologica (New York)* 38:137-144.
- . 1953b. Two new subspecies of *Graptemys pseudogeographica*. *Occ. Pap. Mus. Zool. Univ. Michigan* (546):1-17.
- . 1954. Two new species of the genus *Graptemys*. *Tulane Stud. Zool.* 1:167-186.
- Cahn, A.R. 1937. The turtles of Illinois. *Illinois Biol. Monogr.* 35:1-218.
- Carr, A.F., Jr. 1952. Handbook of turtles. The turtles of the United States, Canada, and Baja California. Cornell Univ. Press, Ithaca, New York.
- and L.T. Marchand. 1942. A new turtle from the Chipola River, Florida. *Proc. New England Zool. Club* 20:95-100.
- Conant, R. and J.T. Collins. 1991. A field guide to reptiles and amphibians: eastern and central North America. Houghton Mifflin Co., Boston.
- Cope, E.D. 1875. Check-list of North American Reptilia and Batrachia. *Bull. U.S. Natl. Mus.* (1):1-104.
- Dobie, J.L. 1981. The taxonomic relationship between *Malaclemys* Gray, 1844 and *Graptemys* Agassiz, 1857 (Testudines, Emydidae). *Tulane Stud. Zool. Bot.* 23:85-102.
- Dundee, H.A. and D.A. Rossman. 1989. The amphibians and reptiles of Louisiana. Louisiana State Univ. Press, Baton Rouge.
- Ernst, C.H. and R.W. Barbour. 1972. Turtles of the United States. The Univ. Press of Kentucky, Lexington.
- and —. 1989. Turtles of the world. Smithsonian Inst. Press, Washington, D. C.
- Folkerts, G.W. and R.H. Mount. 1969. A new subspecies of the turtle *Graptemys nigrinoda* Cagle. *Copeia* 1969:677-682.
- Gaffney, E.S. 1979. Comparative cranial morphology of recent and fossil turtles. *Bull. Amer. Mus. Nat. Hist.* 164:67-376.
- Gibbons, J.W. and J.E. Lovich. 1990. Sexual dimorphism in turtles with emphasis on the slider turtle (*Trachemys scripta*). *Herpetol. Monogr.* 4:1-29.
- Haynes, D. and R.R. McKown. 1974. A new species of map turtle (genus *Graptemys*) from the Guadalupe River system in Texas. *Tulane Stud. Zool. Bot.* 18:143-152.
- Holman, J.A. 1988. The status of Michigan's Pleistocene herpetofauna. *Michigan Acad.* 20:125-132.
- Iverson, J.B. 1992. A revised checklist with distribution maps of the turtles of the world. Privately printed, Richmond, Indiana.
- Jackson, D.R. 1975. A Pleistocene *Graptemys* (Reptilia: Testudines) from the Santa Fe River of Florida. *Herpetologica* 31:213-219.
- Killebrew, F.C. 1977. Mitotic chromosomes of turtles. IV. The Emydidae. *Texas J. Sci.* 29:245-253.
- Lamb, T., C. Lydeard, R.B. Walker, and J.W. Gibbons. 1994. Molecular systematics of map turtles (*Graptemys*): a comparison of mitochondrial restriction site versus sequence data. *Syst. Biol.* 43:543-559.
- Le Sueur, C.A. 1817. An account of an American species of tortoise, not noticed in the systems. *J. Acad. Nat. Sci. Philadelphia* 1:86-88.
- Lovich, J.E. and C.J. McCoy. 1992. Review of the *Graptemys pulchra* group (Reptilia: Testudines: Emydidae), with descriptions of two new species. *Ann. Carnegie Mus.* 61:293-315.
- McCoy, C.J. and R.C. Vogt. 1990. *Graptemys geographica*. *Cat. Amer. Amphib. Rept.* 484:1-484.4.
- McDowell, S.B. 1964. Partition of the genus *Clemmys* and related problems in the taxonomy of the aquatic Testudinidae. *Proc. Zool. Soc. London* 143:239-279.
- McKown, R.R. 1972. Phylogenetic relationships within the turtle genera *Graptemys* and *Malaclemys*. Ph.D. Diss., Univ. Texas, Austin.
- Mount, R.H. 1975. The reptiles and amphibians of Alabama. Auburn Univ. Agr. Exp. Stat., Auburn, Alabama.
- Preston, R.E. 1979. Late Pleistocene cold-blooded vertebrate faunas from the mid-continental United States. I. Reptilia; Testudines, Crocodilia. *Mus. Paleo. Univ. Michigan, Pap. Paleo.* (19):1-53.
- Seidel, M.E. and D.R. Jackson. 1990. Evolution and fossil relationships of slider turtles, p. 68-73. *In* J.W. Gibbons (ed.), Life history and ecology of the slider turtle. Smithsonian Inst. Press, Washington, D.C.
- Shealey, R.M. 1976. The natural history of the Alabama map turtle, *Graptemys pulchra* Baur, in Alabama. *Bull. Florida St. Mus., Biol. Sci.* 21:47-111.
- Slaughter, B.H., W.W. Crook, Jr., R.K. Harris, D.C. Allen, and M. Seifert. 1962. The Hill-Schuler local faunas of the upper Trinity River, Dallas and Denton counties, Texas. *Univ. Texas Bur. Econ. Geol., Rept. Invest.* (48):1-75.
- Smith, H.M. and O. Sanders. 1952. Distributional data on Texan amphibians and reptiles. *Texas J. Sci.* 4:204-219.
- Stejneger, L. and T. Barbour. 1917. A check list of North American amphibians and reptiles. Harvard Univ. Press, Cambridge, Massachusetts.
- Stock, A.D. 1972. Karyological relationships in turtles (Reptilia: Chelonina). *Can. J. Genet. Cytol.* 14:859-868.
- Stovall, J.W. and W.N. McNulty. 1950. The vertebrate fauna and geologic age of Trinity River terraces in Henderson County, Texas. *Amer. Midland Nat.* 44:211-250.
- Vogt, R.C. 1981. Natural history of amphibians and reptiles of Wisconsin. Milwaukee Public Mus., Wisconsin.
- . 1993. Systematics of the false map turtles (*Graptemys pseudogeographica* complex: Reptilia, Testudines, Emydidae). *Ann. Carnegie Mus.* 62:1-46.
- Wilson, R.L. and G.R. Zug. 1966. A fossil map turtle (*Graptemys pseudogeographica*) from central Michigan. *Copeia* 1966:368-369.
- Wood, R.C. 1977. Evolution of the emydine turtles *Graptemys* and *Malaclemys* (Reptilia, Testudines, Emydidae). *J. Herpetol.* 11:415-421.

**C.J. McCoy** (deceased), Carnegie Museum of Natural History, Pittsburgh, PA 15213 U.S.A., and **Richard C. Vogt**, Estación de Biología "Los Tuxtlas," UNAM, Apartado Postal 94, San Andrés Tuxtla, Veracruz, México C.P. 95700.

Primary editor for this account, Michael E. Seidel.

Published 30 December 1994 and Copyright © 1994 by the Society for the Study of Amphibians and Reptiles.